arXiv:1201.3436v2 [cond-mat.mes-hall] 14 Mar 2012

Comment on "Fock-Darwin States of Dirac Electrons in Graphene-Based Artificial Atoms"

The authors of Ref.[1] have computed eigenenergies and eigenstates of a graphene dot in the presence of two-dimensional (2d) harmonic potential and a magnetic field by considering only basis states with energies larger than or equal to zero. We show that the wavefunctions of the nearly chiral Landau level (LL) obtained using this approximate method are inaccurate, and violate a Hellman-Feynman theorem. Corrections lead to significant changes of optical strengths involving nearly chiral states. Moreover, additional basis states lead to new energy levels in the energy spectrum.

In Ref.[1] the following type of basis states of the Hamiltonian matrix are used with energies larger than or equal to zero

$$\psi_{n,m}(\vec{r}) = D_n \begin{pmatrix} -\mathrm{sgn}(n) i \phi_{|n|-1,m}(\vec{r}) \\ \phi_{|n|,m}(\vec{r}) \end{pmatrix}. \qquad (1)$$

The energies of these states are $E_n = \mathrm{sgn}(n)\frac{\hbar v_F}{\ell}\sqrt{2|n|}$ (v_F is the Dirac velocity and ℓ is the magnetic length). First order perturbation theory gives the perturbed wavefunctions of nearly chiral LL states with $n = 0$: $|\psi^{(1)}_{0,m}\rangle \approx |\psi_{0,m}\rangle + C_1|\psi_{1,m+1}\rangle + C_{-1}|\psi_{-1,m+1}\rangle$. These states have all the same angular momentum $J = -m - 1/2$. We find, using $C_{-1} = -C_1$, that the expectation value of the azimuthal velocity $\langle v_\theta\rangle$ is *two times* bigger when both positive and negative energy basis states are included than when only positive energy basis states are included in the perturbation series: $\langle \psi^{(1)}_{0,m}|v_\theta|\psi^{(1)}_{0,m}\rangle = 2\sqrt{2}v_F Re\Big[C_1\langle\phi_{0,m}|e^{i\theta}|\phi_{0,m+1}\rangle\Big]$. Our diagonalization of the Hamiltonian matrix confirms the simple perturbative result above. Fig.1(a) shows that the difference in $\langle v_\theta\rangle$ is about a factor of two.

The azimuthal velocity $\langle v_\theta\rangle$ of states in a quantum dot with $V(r) = \frac{1}{2}\kappa r^2$ should approach the velocity along a Hall bar[2] in the 1d harmonic potential $\frac{1}{2}\kappa x^2$. It is easy to show that the graphene edge electrons of a Hall bar obey the Hellman-Feynman theorem $\frac{1}{\hbar}\frac{\partial \epsilon_n(k)}{\partial k} = \langle\psi_{n,k}|v_y|\psi_{n,k}\rangle$, where the y-component of wavevector is denoted by k and the energy of the n'th LL by $\epsilon_n(k)$. Fig.1(a) displays the values of $\langle v_\theta\rangle$ for nearly chiral states of a dot, and we see that they converge to $\frac{1}{\hbar}\frac{\partial\epsilon_0(k)}{\partial k}$ of the Hall bar (the relevant quantum numbers of the two systems are related by $k\ell = \sqrt{2(m+1)}$[2]). When negative energy basis states are excluded in the Hamiltonian matrix the value of the azimuthal velocity $\langle v_\theta\rangle$ is reduced significantly while the value $\frac{1}{\hbar}\frac{\partial\epsilon_0(k)}{\partial k}$ does not change as much. The Hellman-Feynman theorem is thus violated in this case.

Moreover, numerous new states appear in the energy spectrum, for example, between the energies of $(0,0)$ and $(0,1)$, as shown in Fig.1(b). Note also there are *negative* energy levels, which are absent in Ref.[1]. Some of these new levels give rise to additional optical transitions with significant strength. The appearance of additional levels with positive energies is due to negative energy basis states that get perturbed by the parabolic potential with their energies shifted upward. In addition, we find that the strength of the optical transition $(0,0) \rightarrow (0,1)$ at B=2T with $\alpha = 0.925$ and polarization along x-axis, shown in Fig.3 of Chen et al.[1], is about 2.6 times larger than the correct value obtained by including both positive and negative energy basis states. Physical origin of this difference is anticrossing of $(0,0)$ with other levels in the interval $B < 4$T, see Fig.1(b). The states $(1,0)$ and $(0,1)$ also anticross below 2T with new levels arising from the presence of negative energy basis states. This effect is missing in Ref.[1], and leads to additional modifications of optical strength.



FIG. 1: (a) Velocity $\langle v\rangle$ of the eigenstate with $n = 0$. Here $\alpha = (\kappa\ell^2)/(\frac{\hbar v_F}{\ell}) = 0.2$. Values of velocities using 21 basis states: $\frac{\langle v_\theta\rangle}{v_F}$ is for 2d harmonic potential (circles), and $\frac{1}{\hbar}\frac{\partial\epsilon_0(k)}{\partial k}\frac{1}{v_F}$ are for 1d and 2d harmonic potential systems (triangles). $\frac{\langle v_\theta\rangle}{v_F}$ is for 2d harmonic potential computed excluding negative energy basis states (squares). 11 basis states are included. (b) Energy levels in the Hilbert subspaces with angular momentum $J = -1/2$ (filled circles) and $J = 1/2$ (open circles) are shown together. The number basis states used is 30. Allowed optical transitions obey ΔJ = ±1.

Acknowledgment. This work was supported by the Korean Research Foundation Grant funded by the Korean Government (KRF-2009-0074470).

P.S. Park, S.C. Kim, and S. -R. Eric Yang*
Physics Department, Korea University, Seoul Korea
*eyang812@gmail.com

---

[1] H.-Y Chen, V. Apalkov, and T. Chakraborty, Phys. Rev. Lett. **98**, 186803 (2007).
[2] S. -R. Eric Yang, S. Mitra, A. H. MacDonald and M. P. A. Fisher, J. of the Korean Phys. Soc. 29, S10 (1996).